



Kendrick Regional Center

*Surgical expertise you can trust.*

NAME: \_\_\_\_\_

DATE OF PROCEDURE: \_\_\_\_\_

ARRIVE AT ADMITTING DEPT: \_\_\_\_\_  A.M.  P.M.

PROCEDURE: \_\_\_\_\_

- REPORT TO:**
- ADMITTING DEPT. ST. FRANCIS MOORESVILLE – YELLOW PARKING LOT
  - MOORESVILLE SURGERY CENTER – ST. FRANCIS MOORESVILLE  
GREEN PARKING LOT - 1<sup>ST</sup> FLOOR – SUITE 100
  - MOORESVILLE ENDOSCOPY CENTER – ST. FRANCIS MOORESVILLE  
GREEN PARKING LOT - 1<sup>ST</sup> FLOOR – SUITE 101
  - ADMITTING DEPARTMENT AT ST. FRANCIS BEECH GROVE
  - FRANCISCAN SURGERY CENTER AT ST. FRANCIS SOUTH CAMPUS
  - COMMUNITY HOSPITAL SOUTH – ADMITTING DEPARTMENT

**\*\*BRING SOMEONE WITH YOU TO DRIVE YOU HOME\*\***

***\*FOR CANCELLATION OF THIS PROCEDURE, AT LEAST 24 HOUR NOTIFICATION IS REQUIRED\****

**PREP:** HAVE PRESCRIPTION FILLED THAT IS ATTACHED TO TOP OF THIS PAPER.

**DIET:** THE DAY BEFORE YOUR PROCEDURE YOU WILL HAVE LIQUIDS ONLY.

LIQUIDS: BROTH, TEA, COFFEE, JELL-O, SOFT DRINKS, POPSICLES,  
CLEAR JUICES (APPLE JUICE), SPORTS DRINKS  
**\*NO RED COLORING OR DAIRY PRODUCTS\***

**\*\*THE MOVIPREP CARTON CONTAINS 4 POUCHES AND A DISPOSABLE CONTAINER FOR MIXING. YOU MUST COMPLETE THE ENTIRE PREP TO ENSURE THE MOST EFFECTIVE CLEANSING.\*\***

**FIRST DOSE:** 5:00 pm: EMPTY 1 POUCH A AND 1 POUCH B INTO THE DISPOSABLE CONTAINER. ADD LUKEWARM DRINKING WATER TO THE TOP LINE OF THE CONTAINER. MIX TO DISSOLVE. THE MOVIPREP CONTAINER IS DIVIDED BY 4 MARKS. EVERY 15 MINUTES, DRINK THE SOLUTION DOWN TO THE NEXT MARK (APPROX. 8 OUNCES), UNTIL THE FULL LITER IS COMPLETE. DRINK 16 OZ OF THE CLEAR LIQUID OF YOUR CHOICE.

**SECOND DOSE:** 8:00 pm: REPEAT, MIXING REMAINING POUCH A AND POUCH B IN DISPOSABLE CONTAINER WITH LUKEWARM WATER. DRINK THE SOLUTION EVERY 15 MINUTES UNTIL YOU FINISH THE FULL LITER. DRINK 16 OZ OF THE CLEAR LIQUID OF YOUR CHOICE. YOU MAY CONTINUE TO DRINK CLEAR LIQUIDS UNTIL MIDNIGHT.

**MORNING OF PROCEDURE:**

**NOTHING TO EAT OR DRINK. YOU WILL NOT TAKE ANY MEDICATIONS.** YOU MUST STOP TAKING ASPIRIN, IBUPROFEN, PLAVIX AND PLETAL FOR 7 DAYS PRIOR TO PROCEDURE. YOU MUST STOP TAKING ALL HERBAL SUPPLEMENTS, ROOTS AND VITAMINS 2 WEEKS PRIOR TO SURGERY/PROCEDURE.